

Exploring the costs and outcomes of sexually transmitted infection (STI) screening interventions targeting men in football club settings: preliminary cost-consequence analysis of the SPORTSMART pilot randomised controlled trial

Article (Published Version)

Jackson, Louise J, Roberts, Tracy E, Fuller, Sebastian S, Sutcliffe, Lorna J, Saunders, John M, Copas, Andrew J, Mercer, Catherine H, Cassell, Jackie and Estcourt, Claudia S (2014) Exploring the costs and outcomes of sexually transmitted infection (STI) screening interventions targeting men in football club settings: preliminary cost-consequence analysis of the SPORTSMART pilot randomised controlled trial. *Sexually Transmitted Infections*, 91 (2). pp. 100-105. ISSN 1368-4973

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/53486/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.



OPEN ACCESS

ORIGINAL ARTICLE

Exploring the costs and outcomes of sexually transmitted infection (STI) screening interventions targeting men in football club settings: preliminary cost-consequence analysis of the *SPORTSMART* pilot randomised controlled trial

Louise J Jackson,¹ Tracy E Roberts,¹ Sebastian S Fuller,² Lorna J Sutcliffe,² John M Saunders, Andrew J Copas,^{2,3} Catherine H Mercer,³ Jackie A Cassell,⁴ Claudia S Estcourt²

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/sextrans-2014-051715>).

For numbered affiliations see end of article.

Correspondence to

Professor Tracy Roberts, Health Economics Unit, School of Health and Population Sciences, The Public Health Building, University of Birmingham, Birmingham B15 2TT, UK; t.e.roberts@bham.ac.uk

Received 3 June 2014

Revised 16 September 2014

Accepted 26 October 2014

Published Online First

15 December 2014



Open Access
Scan to access more
free content



► <http://dx.doi.org/10.1136/sextrans-2014-051719>



CrossMark

To cite: Jackson LJ, Roberts TE, Fuller SS, *et al.* *Sex Transm Infect* 2015;**91**:100–105.

ABSTRACT

Background The objective of this study was to compare the costs and outcomes of two sexually transmitted infection (STI) screening interventions targeted at men in football club settings in England, including screening promoted by team captains.

Methods A comparison of costs and outcomes was undertaken alongside a pilot cluster randomised control trial involving three trial arms: (1) captain-led and poster STI screening promotion; (2) sexual health advisor-led and poster STI screening promotion and (3) poster-only STI screening promotion (control/comparator). For all study arms, resource use and cost data were collected prospectively.

Results There was considerable variation in uptake rates between clubs, but results were broadly comparable across study arms with 50% of men accepting the screening offer in the captain-led arm, 67% in the sexual health advisor-led arm and 61% in the poster-only control arm. The overall costs associated with the intervention arms were similar. The average cost per player tested was comparable, with the average cost per player tested for the captain-led promotion estimated to be £88.99 compared with £88.33 for the sexual health advisor-led promotion and £81.87 for the poster-only (control) arm.

Conclusions Costs and outcomes were similar across intervention arms. The target sample size was not achieved, and we found a greater than anticipated variability between clubs in the acceptability of screening, which limited our ability to estimate acceptability for intervention arms. Further evidence is needed about the public health benefits associated with screening interventions in non-clinical settings so that their cost-effectiveness can be fully evaluated.

INTRODUCTION

Young people have the highest risk for sexually transmitted infections (STIs) in the UK, and attempts to engage young men in effective screening have proven to be particularly challenging. Public Health England estimated that 16% of young men aged 15–24 years were screened for chlamydia during 2012 compared with 35% of

young women (assuming one test per person).¹

A range of strategies have been proposed to increase screening participation among men, including outreach in non-clinical settings such as sports venues.² Recent research has suggested that people without a healthcare background can successfully promote certain health behaviours among their peers.³ Among young men in England, football has the highest levels of participation for a team sport.⁴ The *SPORTSMART* pilot trial was designed to develop and evaluate the feasibility and acceptability of two replicable models for promoting STI screening in football clubs (specifically, *Chlamydia trachomatis* and *Neisseria gonorrhoeae* screening), including screening promoted by team captains.

The success of any new intervention in increasing screening uptake needs to be balanced against the resources required to achieve the desired outcome, and additional costs must be evaluated in terms of any additional benefits that can be attributed to them. The objectives of this economic analysis were to obtain cost and outcome data for the alternative screening interventions developed by the *SPORTSMART* pilot trial and to use these data in a preliminary economic evaluation.

METHODS**Pilot trial**

The methods and results of the pilot trial are reported in detail elsewhere.⁵ In brief, a cluster randomised control trial (RCT) design was used involving the allocation of clubs, each with two teams, to one of three trial arms: (1) captain-led and poster STI screening promotion arm; (2) sexual health advisor-led and poster STI screening promotion arm and (3) poster-only STI screening promotion control/comparator arm. The participants were men aged 18 years and over within six amateur football clubs in London. Eligible football clubs were grouped by similar characteristics into three pairs, and then each of the pairs was randomised to a study arm.

The interventions were delivered during the pre-match team briefing. For the captain-led promotion,

the captain delivered a one-time standardised promotion talk of <5 min duration, handed each player a specially designed football-themed sample collection kit and answered any questions from players. In the sexual health advisor-led arm, the standardised promotion talk was delivered by a sexual health advisor from the study clinic. In the poster-only arm, specially designed posters were displayed and kits were made readily available, but no verbal information was given. If men chose to participate, they completed a sample collection kit and placed the completed kit in a secure collection box at the club within an hour after the match ended or they had the option of participating at a later date and posting their sample back to clinic in a discrete postage-paid envelope. Provision of test results and appropriate clinical follow-up was undertaken by the clinical team in the study clinic (off-site), according to routine clinical practice. This included notification of provision of test results via a text message (SMS) by clinic staff.

The primary outcome was the proportion of eligible men accepting the screening offer. The target was to achieve a sample size of 200 men as this would allow the overall acceptance rate to be estimated within 7% if the rate was 50% (ie, a 95% CI 43% to 57%) and within 5% if the rate was higher or lower, assuming minimal variability between clubs.

Economic analysis: overview

For all arms of the trial, resource use data were collected prospectively and unit costs were applied. There were several elements that were common to all of the intervention arms; these related to the recruitment and briefing of the clubs, the materials used within the interventions, and the collection and processing of completed samples. The time taken to recruit, brief and prepare the clubs for the screening intervention was recorded and staffing costs estimated using the Unit Costs of Health and Social Care 2013.⁶ Costs associated with travel to the clubs to prepare for the intervention were also included.

Specially designed posters, football-themed sample kit packs and themed collection containers were used at all locations to promote screening and provide the equipment necessary for participation. We assumed that all these elements were an essential part of the intervention and included their costs in our estimates. We assumed that designs and logos could be reused over a period of 3 years until they became outdated, and thus annuitised all design and editing costs for 3 years at an interest rate of 3%.⁷ We assumed an even number of test kits across intervention arms as data recorded in the trial demonstrated that similar quantities of testing materials were used across arms, as unused items were reused in other clubs. Transport costs associated with returning the completed player samples to the study clinic (off-site) were also included. Although players were given the option to return their samples by post, only one sample was returned by this method, and for the base case, we assumed that the secure collection box would be the only method of specimen return provided.

Additional facilities were required for storage of the samples before they were sent to the laboratory for processing. We assumed that such additional storage would be needed if the intervention were rolled out and that these facilities could be reused over a period of 3 years; we included annuitised costs accordingly. Costs associated with processing samples were estimated using the cross charge between the processing laboratory and the study clinic. Staff time associated with patient administration at the clinic was recorded and costs estimated. We also included costs associated with notifying players about their results and any further costs associated with patient consultation

and treatment. For all trial arms, data were collected on direct health service costs and some of the private costs incurred by the players and captains. The main analysis was conducted from the perspective of the health service (National Health Service).

Resource use and cost definition

Captain- and poster-promoted screening

The estimated cost of the captain-led screening intervention included costs for a member of staff (a healthcare assistant) from the clinic undertaking the sample processing and notification to be on site before and after the intervention to deliver and prepare all the materials and to facilitate the safe return of the completed samples to the clinic, based on recorded practice within the pilot trial and clinical governance requirements. For the base case, it was assumed that the time taken by the team captain to prepare for and deliver the intervention was forgone leisure time and would not impact on health service costs. However, the effect of including these costs or some kind of financial incentive for the captain was analysed as part of the sensitivity analysis.

Sexual health advisor- and poster-promoted screening

The estimated costs for the health advisor-promoted intervention included costs for a sexual health advisor to lead the screening promotion. We assumed that the health advisor would also take the materials to the club, prepare the promotion and ensure the safe return of completed specimen samples to the clinic, in accordance with trial processes and clinical governance requirements and hence included time and travel costs in our estimates.

Poster-promoted screening only (control)

As for the captain-led arm, we assumed that a member of staff (a healthcare assistant) from the clinic undertaking the testing and notification would need to be on site before and after the promotion and included costs accordingly.

Analysis

We conducted a cost-consequences analysis that involves comparing the costs and outcomes associated with all three interventions separately.⁸ This kind of analysis is more appropriate than a full economic evaluation because the current study is a pilot only and a full RCT has not been carried out.

We assessed costs and outcomes in a disaggregated manner for each intervention arm to establish whether any showed clear dominance. Dominance is judged to have occurred when one intervention costs less but is more effective, in terms of the outcome achieved, compared with a different intervention. Conversely, an intervention is dominated if it costs more but is less effective than the comparator. We examined costs and consequences for all three arms. The main analysis is based on the outcome of whether the player accepted the offer of screening. All cost data reported are presented in British pounds in 2012/2013 prices.

A series of one-way deterministic sensitivity analyses were carried out. Uncertainties around all key cost and outcome parameters were analysed, and plausible ranges were specified using information from the trial and from the literature. These analyses included (a) reducing club recruitment time substantially to 4 h per club by developing a higher-level agreement with the Football Association; (b) including an incentive of £1000 for each club to help maximise participation (to reflect practice within the study); (c) including broader societal costs associated with captain participation in screening, with the assumption

Health services research

that the time taken to participate in the intervention was forgone leisure time, valued at 40% of the median hourly wage;^{9 10} (d) reducing intervention costs for the poster-only control arm to analyse the impacts of different staffing arrangements; (e) adjusting the cost of the test kit boxes to account for the logo and design costs associated with unused boxes; (f) increasing sample processing costs and (g) varying uptake levels by study arm. Further sensitivity analyses were carried out but are not reported.

RESULTS

Across all three of the trial arms, 153 men received the intervention and 90 of them accepted the screening offer (59%, 95% CI 35% to 79%, using a robust SE to acknowledge the clustering of participants by club). There was considerable variation in the uptake rates between individual clubs, but results were broadly comparable across study arms (table 1). For the captain-led arm, 56 men received the promotion and 28 accepted (50%). For the health advisor-led arm, 46 men attended the promotion and 31 accepted (67%). For the poster-only control arm, 51 men received the promotion and 31 accepted (61%). There were no positive test results for chlamydia or gonorrhoea in any of the study arms.

Full costs for each of the intervention arms are shown in table 2, and further details are included in online supplementary appendix 1. The results of the pilot trial suggested that total costs were similar across all of the intervention arms, with the total costs of the captain-promoted screening intervention estimated to be £2491.61 compared with £2738.09 for the health advisor-led arm and £2538.09 for the poster-only arm. Overall costs were similar because the highest proportion of costs related to fixed costs, such as staff time for recruiting and briefing the clubs and the equipment required for delivering the promotion.

For all three intervention arms, costs were compared with the main outcome of screening uptake (table 3). The average cost per player tested was comparable across the trial arms using the base case results, with the average cost per player tested for the captain-led promotion estimated to be £88.99 compared with £88.33 for the sexual health advisor-led screening promotion and £81.87 for the poster-only (control) arm.

Sensitivity analysis

As demonstrated in table 4, the results were as follows: (a) decreasing the time needed for club recruitment reduced overall

costs, with the cost per player screened ranging from £60.14 to £66.59; (b) including an incentive in our analysis increased overall costs for all trial arms; (c) including costs for team captains to deliver the promotion made the captain-led arm slightly more expensive, however, as process evaluation revealed that team captains had also informally promoted the screening intervention in other trial arms including these costs for the captain-led arm alone may not be justified; (d) reducing intervention costs for the poster control arm led to a reduction in overall costs for this arm; (e) increasing the costs associated with the test kit boxes (to adjust for costs associated with unused boxes) increased total costs for all intervention arms with estimates per player screened ranging from £84.26 to £91.63; (f) increasing sample processing costs increased costs for all trial arms and (g) varying uptake levels had an effect on the result, emphasising the importance of an accurate estimate of effectiveness.

DISCUSSION

This was an exploratory economic evaluation comparing the costs and outcomes of alternative models for promoting screening among young men. The results as a whole suggest that all these methods of screening promotion are acceptable to players within amateur football clubs, with 153 men receiving the intervention and 90 accepting the offer of screening (59%, 95% CI 35% to 79%). The overall costs associated with the intervention arms were similar. The outcome of average cost per player screened was comparable across all arms, with the average cost per player tested for the captain-led promotion estimated to be £88.99 compared with £88.33 for the sexual health advisor-led screening promotion and £81.87 for the poster-only (control) arm. This outcome is affected by the estimate of the number of players accepting screening, and as our ability to estimate uptake for any single intervention arm is limited, drawing conclusions about the relative costs and consequences of the interventions is not justified. No intervention model can be judged to be dominant, and the average cost per player tested can be seen as comparable across all of the study arms.

It might have been expected that the costs associated with the poster-only control and captain-led arms would be lower than for the health advisor-led arm. However, costs were found to be similar due to the need for a member of clinic staff to be on site and ensure the return of samples to clinic, to meet clinical guidelines. In the event of a rollout of the trial, a satisfactory alternative to this arrangement might be found.

This study has several limitations. It was difficult to draw firm conclusions about the relative cost-effectiveness of the interventions in the trial as screening uptake could not be estimated with precision for any single intervention arm. Two clubs were randomised to each intervention arm; however, we found a greater than anticipated variability between clubs in the acceptability of screening, which limited our ability to estimate acceptability for intervention arms. In addition, the overall target sample size was not achieved due to difficulties in recruitment linked to poor weather and rescheduled matches. We did not capture additional downstream testing that may have occurred as a result of the intervention, and so uptake of STI testing might have been underestimated. As the trial did not identify any positive cases of chlamydia or gonorrhoea, it was not possible to estimate a cost per case diagnosed.

Within the economic analysis, some assumptions were made about how the interventions would operate if they were rolled out, and although these were examined within a sensitivity analysis, they would need to be tested at a larger scale. A one-way

Table 1 Screening uptake for clubs within the study arms

Study arm	Club	Players in changing room	Completed kits returned	Percentage of return
Captain-led	A	26	10	38.5
Captain-led	B	30	18	60.0
		56	28	50.0
Health advisor-led	A	24	10	41.7
Health advisor-led	B	22	21	95.5
		46	31	67.4
Poster-only (control)	A	24	20	83.3
Poster-only (control)	B	27	11	40.7
		51	31	60.8
Total	All	153	90	58.8

Table 2 Health service costs for intervention arms (two clubs per arm)

Resources used	Cost item	Unit cost £*	N	Total cost £*
Intervention costs				
Recruitment of club	Per club	516.88	2	1033.75
Poster pack†	Per pack	53.92	2	107.85
Test kit‡	Per player	5.66	46	260.36
Promotion	Per club	Captain-led: 125.00 Health advisor-led: 225.00 Poster-only: 125.00	2	Captain-led: 250.00 Health advisor-led: 450.00 Poster-only: 250.00
Specimen collection box†	Per club	55.62	2	111.25
Transport of specimen collection box	Per club	135.64	2	271.28
Processing costs				
Additional storage facilities‡	Per club	11.63	2	23.26
Sample processing	Per player tested	10.79	Captain-led: 28 Health advisor-led: 31 Poster-only: 31	302.12 334.49 334.49
Patient admin and notification of results	Per player tested	4.71	Captain-led: 28 Health advisor-led: 31 Poster-only: 31	131.74 145.86 145.86
Total cost				Captain-led: 2491.61 Health advisor-led: 2738.09 Poster-only: 2538.09

*Costs are UK£ (2012/2013).

†Includes costs for the first year of the design elements of the posters, test kit box, pens and specimen collection boxes, annuitised at 3% for 3 years.

‡Includes costs for the first year of the storage facilities, annuitised at 3% for 3 years.

deterministic sensitivity analysis was carried out since this is a preliminary economic analysis alongside a pilot trial and a full probabilistic sensitivity analysis would not be appropriate because of the small sample size and the heterogeneity of uptake rates at club level. Further uncertainties around cost and outcomes parameters would need to be analysed if a full RCT was conducted. Finally, it was intended that players in the control arm would be *uninfluenced* by team captains, but their enthusiasm for the intervention meant that this was not possible, and captains encouraged players to participate in screening via regular team information emails.

The strength of this study is that detailed data on costs and resource use were collected which can inform similar interventions in this area and enable comparisons with other research findings. In addition, data were collected on the number of players who attended the screening promotion events, and thus it is possible to estimate the level of uptake of screening. Often with such health promotion interventions, the number of people exposed to a particular intervention is unknown.

Very little information exists about the cost-effectiveness of screening programmes in non-clinical settings. A recent systematic review of studies concerned with chlamydia and gonorrhoea screening outreach programmes only identified three with information on costs.¹¹ Buhner-Skinner *et al*¹² calculated a cost per test carried out for outreach clinics in Australia but did not

include staff time, transport and setup costs. Morris *et al* calculated the costs associated with two Californian youth programmes but did not include additional time associated with volunteer input.¹³ Detailed costings were also provided by a study evaluating the cost-effectiveness of a multifaceted community intervention to increase screening in Stockholm.¹⁴ However, due to the nature of the intervention, which involved a large-scale publicity campaign and expanded access to testing facilities, it is difficult to compare these results with the SPORTSMART study. The findings of a costing study of chlamydia screening within primary care suggest that the costs per case screened for the SPORTSMART study are higher than would usually be expected in a UK primary care setting.¹⁵ However, the additional public health benefits associated with outreach activities would also need to be taken into account and achieving adequate coverage of screening in primary care, especially in men, is challenging.

Conclusions

This preliminary economic evaluation has shown that similar costs and outcomes were demonstrated across all three study arms. The fact that the control arm was unintentionally 'enhanced' by team captains suggests that they can have an important influence on screening uptake, irrespective of whether they take on a formal role in promoting the intervention and the potential costs

Table 3 Comparison of costs and outcomes for the intervention arms

Intervention arm*	Total cost £†	Number of players tested	Per cent accepting screening offer	Average cost per player screened £†
Captain-led	2491.61	28	50	88.99
Health advisor-led	2738.09	31	67	88.33
Poster-only (control)	2538.09	31	61	81.87

*Includes costs and outcomes for both clubs in each trial arm.†Costs are UK£ (2012/2013).

Table 4 Sensitivity analysis: selected results

	Original value	Revised value	Captain-led arm: total cost* (average cost per player screened)	Health advisor-led arm: total cost* (average cost per player screened)	Poster-only arm: total cost* (average cost per player screened)
Base case					
(a) Reducing club recruitment time to 4 h per club			£2491.61 (£88.99)	£2738.09 (£88.33)	£2538.09 (£81.87)
(b) Including £1000 incentives for each club	£516.88†	£180†	£1817.86 (£64.92)	£2064.34 (£66.59)	£1864.34 (£60.14)
(c) Including costs for team captains to deliver promotion		£1000†	£4491.61 (£160.41)	£4738.09 (£152.84)	£4538.09 (£146.39)
(d) Reducing promotion costs for the poster-only control arm		£2.68†	£2496.87 (£89.17)		
(e) Increasing costs for testing boxes	£125.00†	£50.00†			
(f) Increasing sample processing costs	£5.66†	£7.27†	£2565.67 (£91.63)	£2812.15 (£90.71)	£2388.09 (£77.04)
(g) Varying uptake levels by study arm	£10.79†	£16.19†	£2642.67 (£94.38)	£2905.34 (£93.72)	£2705.34 (£87.27)
	Captain-led: 50.0% Health advisor: led: 67.4% Poster-only: 60.8%	38.5–95.4%	£2246.10–£2818.77 (£124.78–£64.06)	£2446.10–£3018.77 (£135.89–£68.61)	£2277.09–£2896.24 (£113.85–£59.11)

Costs are UK£ (2012/2013).

*Includes costs and outcomes for both clubs in each trial arm.

†Per club, †Per item.

associated with their informal and formal input need to be carefully considered. Further research is needed to explore the public health benefits associated with screening interventions in non-clinical settings so that their cost-effectiveness can be fully evaluated. This would allow the potentially higher costs associated with such 'outreach' activities to be weighed against their wider health benefits so that policymakers can be fully informed about which approaches offer the best value for money.

Key messages

- ▶ Although acceptance rates were highly variable between clubs, levels of uptake were broadly comparable across all study arms.
- ▶ The overall costs associated with the intervention arms were similar.
- ▶ No intervention model was judged to be dominant, and the average cost per player tested was comparable across all of the study arms.
- ▶ Further research is needed to investigate the public health benefits associated with screening interventions in non-clinical settings so that their cost-effectiveness can be fully evaluated.

Author affiliations

¹Health Economics Unit, School of Population and Health Sciences, College of Medical and Dental Sciences, University of Birmingham, Birmingham, UK

²Centre for Immunology & Infectious Disease: Sexual Health & HIV, Blizard Institute, Queen Mary, University of London, Barts and the London School of Medicine and Dentistry, London, UK

³Centre for Sexual Health and HIV Research, Faculty of Population Health Sciences, University College London, London, UK

⁴Division of Primary Care & Public Health, Brighton and Sussex Medical School, University of Brighton, Falmer, Brighton, UK

Handling editor David A Lewis

Acknowledgements The authors would like to thank the following people for their assistance with this study: Mr Simon Morgan, Head of Community Development, Barclay's Premier League; Mr David Higgins; Ms Stephanie Hanson; Mr Timothy Hill; and all the participating clubs and players. We would also like to thank the anonymous referees for their helpful comments on our draft manuscript.

Contributors CSE, IJS, AJC, CHM, JAC and TER are co-applicants on the NIHR BALLSEYE study. CSE was the principal investigator and led the research for the trial. SSF was study researcher and led the conduct of the trial. TER designed the economic evaluation and supervised the analysis. The analysis was carried out by LJJ who wrote the first draft as the lead author. All authors contributed to the design of the study and provided input to this manuscript.

Funding The SPORTSMART study is part of the NIHR-funded BALLSEYE Programme and is led by Dr Claudia Estcourt. This paper is independent research funded by the National Institute for Health Research (BALLSEYE Programme, 'Targeting Men for Better Sexual Health', RP-PG-0707-10208). The views expressed in this publication are those of the authors and not necessarily those of the NHS, the National Institute for Health Research or the Department of Health.

Ethics approval Ethical approval was obtained for the clinical trial from the National Research Ethics Service (reference 13/SC/0029). Ethical approval was not required for the economic evaluation.

Competing interests None.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

- 1 Public Health England. *Health Protection Report (Weekly Report)*, June 2013, 7(23). <http://www.hpa.org.uk/hpr/archives/2013/hpr2313.pdf> (accessed 1 Nov 2013).

- 2 Baraitser P, Blake S on behalf of NHS National Chlamydia Screening Programme. *Involving young men in chlamydia screening: a practical guide*. London: Health Protection Agency, 2009. http://www.chlamydia-screening.nhs.uk/ps/resources/guidelines/NCSP_Involving_young_men.pdf (accessed 1 Nov 2013).
- 3 Williamson LM, Hart GJ, Flowers P, *et al*. The Gay Men's Task Force: the impact of peer education on the sexual health behaviour of homosexual men in Glasgow. *Sex Transm Infect* 2001;77:427–32.
- 4 Saunders JM, Mercer CH, Sutcliffe LJ, *et al*. Where do young men want to access STI screening? A stratified random probability sample survey of young men in Great Britain. *Sex Transm Infect* 2012;88:427–32.
- 5 Fuller SS, Mercer CH, Copas AJ, *et al*. The SPORTSMART study: a pilot randomised controlled trial of sexually transmitted infection (STI) screening interventions targeting men in football club settings. *Sex Transm Infect* 2015;91:106–10.
- 6 Curtis L. *Unit costs of health and social care 2012*. Kent: Personal Social Services Research Unit, 2012.
- 7 Roberts TE, Tsourapas A, Sutcliffe L, *et al*. Is Accelerated Partner Therapy (APT) a cost-effective alternative to routine patient referral partner notification in the UK? Preliminary cost–consequence analysis of an exploratory trial. *Sex Transm Infect* 2012;88:16–20.
- 8 Drummond MF, O'Brien B, Stoddart GL, *et al*. *Methods for the economic evaluation of health care programmes*. Oxford: Oxford University Press, 2005.
- 9 Office of National Statistics. *Annual Survey of Hours and Earnings, 2013 Provisional Results*. 2013. <http://www.ons.gov.uk/ons/rel/ashe/annual-survey-of-hours-and-earnings/2013-provisional-results/stb-ashe-statistical-bulletin-2013.html> (accessed 11 Dec 2013).
- 10 Robinson S, Roberts T, Barton P, *et al*. Healthcare and patient costs of a proactive chlamydia screening programme: the Chlamydia Screening Studies project. *Sex Transm Infect* 2007;83:276–81.
- 11 Hengel B, Jamil MS, Mein JK, *et al*. Outreach for chlamydia and gonorrhoea screening: a systematic review of strategies and outcomes. *BMC Public Health* 2013;13:1040.
- 12 Buhner-Skinner M, Muller R, Menon A, *et al*. Novel approach to an effective community-based chlamydia screening program within the routine operation of a primary healthcare service. *Sex Health* 2009;6:51–6.
- 13 Morris S, Bauer H, Chartier M, *et al*. Relative efficiency of chlamydia screening in non-clinical settings in two California counties. *Int J STD AIDS* 2010;21:52–6.
- 14 Deogan CL, Bocangel MKH, Wamala SP, *et al*. A cost-effectiveness analysis of the Chlamydia Monday-A community-based intervention to decrease the prevalence of chlamydia in Sweden. *Scand J Public Health* 2010;38:141–50.
- 15 Cassell JA, Estcourt C, Llewellyn C, *et al*. The relative clinical and cost-effectiveness of three contrasting approaches to partner notification for curable sexually transmitted infections (STIs): a cluster randomised trial in primary care. *Health Technol Assess* 2014; In press.

STI

Exploring the costs and outcomes of sexually transmitted infection (STI) screening interventions targeting men in football club settings: preliminary cost-consequence analysis of the *SPORTSMART* pilot randomised controlled trial

Louise J Jackson, Tracy E Roberts, Sebastian S Fuller, Lorna J Sutcliffe, John M Saunders, Andrew J Copas, Catherine H Mercer, Jackie A Cassell and Claudia S Estcourt

Sex Transm Infect 2015 91: 100-105 originally published online December 15, 2014
doi: 10.1136/sextrans-2014-051715

Updated information and services can be found at:
<http://sti.bmj.com/content/91/2/100>

These include:

Supplementary Material

Supplementary material can be found at:
<http://sti.bmj.com/content/suppl/2014/12/15/sextrans-2014-051715.DC1.html>

References

This article cites 8 articles, 6 of which you can access for free at:
<http://sti.bmj.com/content/91/2/100#BIBL>

Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Open access](#) (179)
[Screening \(epidemiology\)](#) (883)
[Screening \(public health\)](#) (883)

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>